

## Clinical Image

Open Access, Volume 4

# Calcified coronary artery aneurysm

**Kaavya Nair<sup>1\*</sup>; Nandan Anavekar<sup>2</sup>**

<sup>1</sup>Research Fellow, Mayo Clinic, Rochester MN, USA.

<sup>2</sup>Program Director for Adult Cardiovascular Medicine, Mayo Clinic, USA.

**\*Corresponding Author: Kaavya Nair**

Research Fellow, Mayo Clinic, Rochester MN, USA.

Phone: 818-451-5294;

Email: kaavyanjali96@gmail.com

Received: Aug 03, 2023

Accepted: Aug 23, 2023

Published: Aug 30, 2023

Archived: www.jcimcr.org

Copyright: © Nair K (2023).

DOI: www.doi.org/10.52768/2766-7820/2567

**Keywords:** Cardiology; Imaging; Decision-making.


### Description

58-year-old woman presented to the hospital with sudden onset right sided abdominal and chest pain. Coronary angiogram demonstrated calcified aneurysms (yellow arrows) at the bifurcation left main which were in communication with one another (green arrow). The distal aneurysm was noted to be partially thrombosed (red arrow) (Figure 1).

Cardiothoracic surgery did not recommend any surgery to coronary arteries, given stability of aneurysm size when compared to prior imaging. Notably, the patient was found to have a large pericardial effusion without signs of tamponade. After pericardiocentesis, there was a marked improvement of symptoms.

**Funding:** None

**Conflicts of interest:** None.



**Figure 1:** Coronary angiogram.