

## Clinical Image

Open Access, Volume 4

# Severe scrotal trauma in polytrauma patients

**Cheikh Saad Bouh Khattraty\***; Mohamed Amine Malki; Mohammed Benamar; Mustapha Ahsaini; Soufiane Mellas;

Jalal Eddine EL Ammari; Mohammed Fadl Tazi; Mohammed Jamal Fassi; Moulay Hassan Farih

Department of Urology, University Hospital Hassan II, Fez Morocco.

**\*Corresponding Author: Bouh Khattraty CS**

Department of Urology, University Hospital Hassan II, Fez Morocco.

Email: dr.khattraty@gmail.com

Received: Nov 20, 2023

Accepted: Dec 14, 2023

Published: Dec 21, 2023

Archived: www.jcimcr.org

Copyright: © Bouh Khattraty CS (2023).

DOI: www.doi.org/10.52768/2766-7820/2748

### Background

Testicular or scrotal trauma is rare, but most often occurs after an assault with a firearm. Most patients have only isolated scrotal or testicular trauma.

However, their incidence is increasing with the rise in road accidents involving motorcycles and bicycles [1].

We report the case of a 28-year-old motorcyclist who was hit by a car, resulting in cerebral, abdominal, traumatological and urological trauma, with both testicles exposed, as shown in the following image.

Both testicles were viable, but the scrotal skin was completely absent, the scrotal lesion extending to the base of the penis (Figure 1).

After resuscitation, the patient was trimmed in the operating room.



Figure 1: Severe scrotal trauma

### References

1. Grigorian A, Livingston JK, Schubl SD, Hasjim BJ, Mayers D, Kuncir E, Barrios C, Joe V, Nahmias J. National analysis of testicular and scrotal trauma in the USA. Res Rep Urol. 2018 Aug 10, 30128306, 10:51-56. doi: 10.2147/RRU.S172848.

**Citation:** Bouh Khattraty CS, Amine Malki M, Benamar M, Ahsaini M, Mellas S, et al. Severe scrotal trauma in polytrauma patients. J Clin Images Med Case Rep. 2023; 4(12): 2748.