

Clinical Image

Open Access, Volume 4

Surprises in surgery: Clinical image

Sergio Castorina^{1,2,3}; Ornella Coco¹; Mariacarla Castorina¹; Stefano Puleo¹

¹Mediterranean Foundation "GB Morgagni," Catania, Italy.

²Department of Medical, Surgical Sciences and Advanced Technology, Catania, Italy.

³Mediterranean Foundation "GB Morgagni," Catania, Italy.

***Corresponding Author: Ornella Coco**

Mediterranean Foundation "GB Morgagni," Catania
Italy.

Tel: 3470091594; Email: ornellacoco19@gmail.com

Received: Nov 25, 2023

Accepted: Dec 22, 2023

Published: Dec 29, 2023

Archived: www.jcimcr.org

Copyright: © coco O (2023).

DOI: www.doi.org/10.52768/2766-7820/2768

Introduction

Gallbladder agenesis represents the 0.01-0.04% of congenital anomalies (0.016 in autopsies). Patient become symptomatic in over the 20% of cases with false-positive imaging of cholelithiasis on ultrasound examination that leads to elective laparoscopic cholecystectomy. Pancreas divisum (PD) is the most frequent congenital disorder of the pancreas occurring in 4-14% of population and cause of recurrent acute pancreatitis. Three types of PD are known: type 1 (70%) total failure of fusion; type 2 (20%) dominant dorsal drain in absence of Wirsung; type 3 (5-10%) inadequate connection between dorsal and ventral branch.

Clinical case

A 53 years old man was admitted for recurrent pain in right hypochondrium and epigastrium, ultrasound (US) findings of gallstones in scleroatrophic gallbladder without blood count values of cholestasis. Laparoscopy showed a normal common biliary duct (CBD) in absence of gallbladder in its normal site and in the most common sites for ectopia (Figure 1). Surgical procedure was suspended and magnetic resonance cholangiopancreatography was utilized according to Meld's algorithm, in order to verify any other malformations. MRCP showed normal

intra ed extrahepatic biliary tree (Figure 2), gallbladder agenesis and a suspect Pancreas divisum (type 3) (Figure 3).

Conclusion

In presence of an incidental diagnosis of gallbladder agenesis it is suggested to avoid further surgical procedures and to follow the Meld's algorithm for confirming gallbladder agenesis by MRCP. Conservative therapy and/or sphincterotomy are the following options on the base of patient's symptoms.




Figure 1:




Figure 2:




Figure 3:

References

1. Sachin Malde. Gallbladder agenesis diagnosed intra-operatively: a case report. Journal of Medical Case Reports. 2010; (4): 285.
2. Sergio Castorina, Roberto Scilletta, Jacques Domergue. Gallbladder agenesis: laparoscopic views of a significant diagnostic challenge. SurgRadiol Anat. 2014; 36: 617-620. DOI: 10.1007/s00276-014-1263-0.
3. Maryselle Winters, Derek T Clar, Kelly Van Fossen. Intraoperative Diagnosis of Gallbladder Agenesis. Am Surg. 2023; 89(6): 2844-2846. doi: 10.1177/00031348211060440.
4. Oktay Yener et al. Agenesis of the gallbladder diagnosed by magnetic resonance cholangiography: report of a case and review of the literature. Prague Med Rep. 2015; 116(1): 52-6. Doi: 10.14712/23362936.2015.4