

**Clinical Image**

Open Access, Volume 5

**Hemifacial spasm and blepharospasm caused by vertebral artery dolichoectasia****Entela Basha<sup>1\*</sup>; Gentian Vyshka<sup>2</sup>; Eris Ranxha<sup>1</sup>; Artur Xhumari<sup>3</sup>**<sup>1</sup>Neurovascular Service, "Mother Teresa" University Medical Center, Tirana, Albania.<sup>2</sup>Biomedical and Experimental Department, University of Medicine in Tirana, Albania.<sup>3</sup>Service of Neurosurgery, "Mother Teresa" University Medical Center, Tirana, Albania.**\*Corresponding Author: Entela Basha**Neurovascular Service, "Mother Teresa" University  
Medical Center, Tirana, Albania.

Email: bashaentela@gmail.com

Received: Mar 12, 2024

Accepted: Apr 03, 2024

Published: Apr 10, 2024

Archived: www.jcimcr.org

Copyright: © Basha E (2024).

DOI: www.doi.org/10.52768/2766-7820/2978

**Abstract**

Hemifacial spasm and blepharospasm are crippling medical conditions that severely affect the quality of life of patients. Among differential diagnosis, the ectasia of vertebro-basilar vessels should be taken into account, and treated accordingly. Authors report the case of a 60-years old male with a two-year history of involuntary intermittent twitching of the right eye and right side of the upper lip. MRI of the brain suggested a dolichoectasia of the vertebral artery. The condition was almost refractory to drugs and the patient was referred for neurosurgical intervention.

**Keywords:** Hemifacial Spasm; Blepharospasm; Neurovascular Conflict.


**Description**

Hemifacial spasm and blepharospasm are subtypes of peripheral movement disorder. They are divided into primary and secondary categories based on their etiology. Primary hemifacial spasm usually follows a neurovascular conflict between the posterior circulation vessels and facial nerve root, while common secondary causes are the Bell's palsy, traumatic injury of the facial nerve, demyelination, and stroke [1,2]. A 60 years old Albanian man reported a two-year history of involuntary intermittent twitching of the right eye and right side of the upper lip that came to his attention only recently due to increased frequency. The patient did not report any other medical illnesses, nor a positive family history of involuntary movements. When asked, he denied contractions in other body parts. Unilateral involuntary contractions affecting his right orbicularis oculi and right orbicularis oris were noted during neurological examination. Spasms were isolated in these two muscles and no other objective sign was found.

Routine biochemical tests and cell blood count were normal. Magnetic Resonance Imaging showed the neurovascular conflict between the right facial nerve and right dolichoectasia of the vertebral artery (Figure 1). The neurosurgical team recommended microvascular decompression surgery of the facial nerve as a definitive treatment.

The patient refused it and preferred pharmacological treatment. We decided to start the treatment with Clonazepam 2 mg per day. The patient came after several months without any clinical improvement. During these follow-up visits, the neurological examination remained unchanged. Treatment with Baclofen 40 mg per day was started, without any improvement. The patient decided to undergo surgery after unsuccessful pharmacological treatment. The spasms resolved the next post-operative day, without recurrence after a one-year follow-up.

Dolichoectasia of the vertebro-basilar system is a normal anatomic variant, where the arteries are enlarged and tortuous. It is usually not associated with neurological complications. Some



**Figure 1:** Left and middle inset: axial MRI images, plain and contrast-phase (angio) showing the dilated, ectatic right vertebral artery in close contact with brainstem structures. Right inset: angio-MRI imaging of the vessel tortuosity, right vertebral artery.

cases have reported manifestations of dolichoectasia of vertebro-basilar system with hydrocephaly, bulbar compression, and vestibulocochlear symptoms [3]. In most cases, primary hemifacial spasms and blepharospasm relate to a neurovascular conflict of the anterior inferior cerebellar artery and posterior inferior cerebellar artery, while our patient's symptoms were caused by dolichoectasia of the vertebral artery. Herewith we emphasize the importance of surgical treatment as a permanent solution to this condition.

### References

1. Campos-Benitez M, Kaufmann AM. Neurovascular compression findings in hemifacial spasm. *J Neurosurg.* 2008; 109(3): 416-20.
2. Yaltho TC, Jankovic J. The many faces of hemifacial spasm: differential diagnosis of unilateral facial spasms. *Mov Disord.* 2011; 26(9): 1582-92.
3. Huh G, Bae YJ, Woo HJ, Park JH, Koo JW, Song JJ. Vestibulocochlear Symptoms Caused by Vertebrobasilar Dolichoectasia. *Clin Exp Otorhinolaryngol.* 2020; 13(2): 123-132.